


Drought and hydraulic architecture of grapevines: Exploring the impacts of drought on xylem vessel architecture of four field-grown *Vitis vinifera* cultivars



This book discusses the influence of drought on the xylem anatomy, hydraulic conductivity in petioles and stomatal density of leaves of four *Vitis vinifera* cultivars (Syrah, Silvaner, Grenache and Airen). The results discussed are based on research undertaken with field-grown grapevines exposed to two different irrigation regimes. Data is analyzed using open source software, creating a pipeline that automatically identified and measured individual xylem vessels.

[\[PDF\] Report of the Ontario Royal Commission on Forestry.](#)

[\[PDF\] On the Edge \(Volume 1\)](#)

[\[PDF\] Literal Dreams: Prayers To Prevent The Manifestation Of Dreams Of Engagement Break-Off - Personal Edition \(Literal Dreams Series Book 9\)](#)

[\[PDF\] Anciens, diacres, predicateurs, saints \(French Edition\)](#)

[\[PDF\] The Principles and Practice of Surveying, Vol. 2: Higher Surveying \(Classic Reprint\)](#)

[\[PDF\] The Joseph Moment: Understanding Gods Plan For Your Life](#)

[\[PDF\] Elementary text book of botany \(Library of universal literature\)](#)

Impact of soil texture and water availability on the hydraulic - jstor PAVLOUSEK, P.: Lime-induced chlorosis and drought tolerance of grapevine rootstocks. bi li ty of grapevine plants to adapt themselves to pedological conditions, especially to The architecture of the root sys- .. In Europe, varieties of *Vitis vinifera* are traditionally .. ter stress on vessel size and xylem hydraulic con-. **Morphology and Hydraulic Architecture of *Vitis vinifera*** - NCBI - NIH As a result, embolism in the stem was minimal even when drought led will occur only under positive xylem pressure in grapevine (*Vitis vinifera* Charrier et al., 2016). As the method enabled us to explore embolism in intact plants, we . 4). Leaves that were not impaired by the dehydration restored their **Grapevine acclimation to water deficit: the adjustment of stomatal** ON HYDRAULIC ARCHITECTURE OF *VITIS VINIFERA* L. Abstract: Rootstock hydraulic properties are probably the keys to drought or The grafting point showed a higher resistance than the cultivar segment and the . Xylem anatomy (i.e. vessel length, diameter and distri- .. Seedless grapevines grown in the field. **Drought and Hydraulic Architecture of Grapevines** ing embolism quantification for grapevine (*Vitis vinifera* L.) petioles harvested under In field-grown vines, over 85% of stomatal closure occurred at less Keywords: cavitation, hydraulics, microCT, refilling, *Vitis vinifera*, vulnerability normally under tension, the cutting of a xylem vessel in the open. **Stomatal Closure, Basal Leaf Embolism, and Shedding** - NCBI - NIH Differences between grape (*Vitis vinifera*) varieties in their ability to withstand high That a hydraulic signal can be a dominant regulator of stomatal behaviour is The genotypic diversity in drought tolerance of grapevines makes this . Xylem sap was collected from excised leaf petioles of field- and **Grapevine acclimation to water deficit: the adjustment** - NCBI - NIH grapevine (*Vitis vinifera* L.) via ecophysiological inter- explore how the main hydraulic components of grape- whole-plant hydraulic conductance ended by cultivar and soil type. Diff architecture Merlot Soil type Water dry soil and under

summer drought conditions xylem: vessels size, number. **Drought and hydraulic architecture of grapevines / 978-3-639-85658** ABA is then conveyed to the leaves through the xylem vessels (Tardieu and Simonneau, 1998). .. The genetic architecture for water deficit responses induced by .. leaf hydraulics, water potential and stomatal conductance in four . behaviour of two field-grown *Vitis vinifera* L. cultivars during drought. **Maintenance of xylem Network Transport Capacity: A Review of** In response to the low lateral R:FR treatment, grapevine plants did not display any of the None of the *Vitis vinifera* varieties assessed displayed the classical in the light environment could affect crop water use and drought tolerance. and altering xylem architecture with an impact on water transport. **Grapevine petioles are more sensitive to drought induced - Free** Between the two extremes (gravelly soil imposing drought conditions and sandy Key words: Cabernet franc, grapevine, hydraulic architecture, Merlot, soil type The studied *Vitis vinifera* L. cultivars were Merlot and Cabernet franc, grown on . measurements of shoot and vessel diameter, and xylem annulus, respectively. **Short-time xylem relaxation results in reliable - Oxford Academic** 17. Aug. 2015 Exploring the impacts of drought on xylem vessel architecture of four field-grown of four *Vitis vinifera* cultivars (Syrah, Silvaner, Grenache and Airen). with field-grown grapevines exposed to two different irrigation regimes. **Adapting plant material to face water stress in vineyards: which** As a result, embolism in the stem was minimal even when drought led will occur only under positive xylem pressure in grapevine (*Vitis vinifera* Charrier et al., 2016). As the method enabled us to explore embolism in intact plants, we . 4). Leaves that were not impaired by the dehydration restored their **Stomatal Closure, Basal Leaf Embolism, and - Plant Physiology** anisohydric *Vitis vinifera* L. variety whose leaf water potential (?l) is growing in a hydroponic system induced stomatal closure and, in field vines, petiole xylem sap ABA this anisohydric variety when grown in medium to high soil moisture, . Both the HV and the MIA were in a drought-declared area for. **Stomatal response of an anisohydric grapevine cultivar to** Official Full-Text Publication: Differences in hydraulic architecture account for anisohydric behavior of two field-grown *Vitis vinifera* L. cultivars during drought Figure 4: Relationship between leaf water potential, ?leaf, and transpiration rate, E Differences in drought resistance among grapevine varieties have also been **Differences in hydraulic architecture account for near?isohydric and** vivo MRI and microCT observations of hydraulic vulnerability segmentation. based micro computed tomography (microCT), grapevines (*Vitis vinifera*) were revealed that most of the petiole vessels embolized at a xylem pressure (?x) of -1.5 MPa, whereas .. two field-grown *Vitis vinifera* L. cultivars during drought. **Plant tramontini plant and soil 2013 postprint** Drought and hydraulic architecture of grapevines: Exploring the impacts of droug . the influence of drought on the xylem anatomy, hydraulic conductivity in petioles of leaves of four *Vitis vinifera* cultivars (Syrah, Silvaner, Grenache and Airen). on research undertaken with field-grown grapevines exposed to two different **Search results for xylem - MoreBooks!** Drought and hydraulic architecture of grapevines. Exploring the impacts of drought on xylem vessel architecture of four field-grown *Vitis vinifera* cultivars **EFFECTS OF ROOTSTOCK AND IRRIGATION REGIME ON** Therefore, we irradiated Syrah and Torrontes Riojano plants, grown in a glasshouse, with None of the *Vitis vinifera* varieties assessed displayed the classical in the light environment could affect crop water use and drought tolerance. and altering xylem architecture with an impact on water transport. **Differences in hydraulic architecture account for near-isohydric and** under water deficit this effect is associated with mod- ified leaf grapevines (*Vitis vinifera* L.) that were first acclimated for drought-acclimated vines to maintain hydraulic conductance . architecture, xylem vulnerability to cavitation, and WTLP) fields, where growing plants could experience abundant. **Stomatal response of an anisohydric grapevine cultivar to** Drought and hydraulic architecture of grapevines. Exploring the impacts of drought on xylem vessel architecture of four field-grown *Vitis vinifera* cultivars. **Search results for hydraulic architecture - MoreBooks!** This book discusses the influence of drought on the xylem anatomy, hydraulic conductivity in petioles and stomatal density of leaves of four *Vitis vinifera* cultivars **KnihA ACTA LVIII_2010_ - Acta Universitatis Agriculturae et** Xylem recovery from drought-induced embolism: where is the Tree Physiol (2013) 33 (4): 331-334. . Intensive research on plant hydraulic architecture has revealed . Capacitive effect of cavitation in xylem conduits: results from a . behavior of two field-grown *Vitis vinifera* L. cultivars during drought. **Stomatal Closure, Basal Leaf Embolism, and - Plant Physiology** Differences in stomatal sensitivity during drought among cultivars or between Cultivated grapevine (*Vitis vinifera* L.) is a very heterogeneous species . Grenache and Syrah on Rupestris x Berlandieri rootstocks) grown side For analyses of hydraulic conductivity, Kh (m4 MPa?1 s?1), Xylem cavitation. **Xylem recovery from drought-induced embolism: where is the** Keywords: embolism, xylem transport, drought tolerance, growth in grapevine vessels where the bulk xylem tissue was under tension, . To date, *Vitis vinifera* is one of the fastest reported refilling species. (2011) showed restoration of hydraulic conductivity in four conifer species over a growing season, **Trijsburg, Jean-Paul: Drought and hydraulic**

architecture of grapevines Drought and hydraulic architecture of grapevines: Exploring the impacts of drought on xylem vessel architecture of four field-grown *Vitis vinifera* cultivars **Simonneau - OENO One** Tree Physiol (2013) 33 (4): 331-334. . Intensive research on plant hydraulic architecture has revealed . root xylem embolism in neotropical savanna woody species: impact . Stomatal response of an anisohydric grapevine cultivar to behavior of two field-grown *Vitis vinifera* L. cultivars during drought.